Land surface Interactions with the Atmosphere over the IberianSemi-arid Environment (LIAISE): 1st modelling intercomparison

M.A. Jiménez¹, J. Cuxart¹, A. Grau¹, (1) Universitat de les Illes Balears, Palma, Spain

A. Boone², S. Donier², P. Le Moigne², (2) CNRM, Meteo-France/CNRS, Toulouse, France


J.R. Miró³, Jordi More³, (3) Servei Meteorològic de Catalunya, Barcelona, Spain

A. Tiesi⁴, P. Malguzzi⁴, (4) CNR-ISAC, Bologna, Italy


J. Brooke⁵ and Martin Best⁵ (5) Met Office, Exeter, UK








The 1st mesoscale intercomparison case





The 1st mesoscale intercomparison case

16-18 July 2016

- * clear skies, A conditions
- * thermal heterogeneities

* locally/basin/mesoscale generated

winds (interact)


Models

MesoNH (MNH)

MOLOCH (MOL)

Unified Model (UM)

WRF



Model setup


- 36h run (from 16 July at 1800 UTC to 18 July at 0600UTC)
- 2 nested domains (1-way): 2km and 400m resolution
- Vertical resolution (2m and stretched above, 85 levels)
- Initial/Lateral BC: ECMWF
- **Differences**: Turbulence, Radiation (5min), Surface


10 m (agl) wind vectors (MesoNH)

17th July 0600 UTC E-wind prevail


17th July 1500 UTC

SB front interacts with local winds







Raïmat (VK) - WEST




Validation using AWS

Moloch WRF MesoNH UM






El Poal (V8) - FLOOD



Validation using AWS

Moloch WRF MesoNH UM



Mean BIAS (model-obs)


Validation using AWS


MesoNH Moloch


UM


ALL run, DAY (1000-1400 UTC), NIGHT (0000-0400 UTC)

Temperature: about +1.5°C
























time (hours)

time (UTC)

Variability of the surface cover



 Models present differences in the surface parameters (LAI, fveg, albedo, ...)

Summary

- The case 16-18 July 2016 is taken for the 1st mesoscale intercomparison.
- Locally-generated circulations
 (interaction between local, basin, mesoscale)
- Results (known features): models are able to reproduce the general patterns of the region <u>BUT</u>:
- Models tend to overestimate wind speed (daytime)
- Difficulties in reproducing nocturnal nearly calm conditions
- Temperatures are overestimated (specially during nighttime)

Summary

- Models are not able to reproduce the heterogeneities:
 Surface model (processes included, irrigation)
 Surface parameters & initialitation (irrigated, rainfed,... zones)
 Parameterizations (turbulence, advection, radiation)
- Sensitivity tests (work in progress)
- ✓ Initial and lateral BC (GFS, NCEP)
- ✓ Surface features

(soil moisture, vegetation, surface model...)

- ✓ Spatial resolution
- After testing models + LIAISE campaign: possible future GEWEX intercomparison

THANKS

session 3. Remote and in situ measurements (27th May 2021)

3P4: Surface thermal heterogeneities in the eastern Ebro basin and their impact on regional circulations, Torres et al.

14h30: Land surface Interactions with the Atmosphere over the Iberian Semiarid Environment (LIAISE) Project: Field Campaign Update, A. Boone et al.

14h45: An overview of the Analysis of Precipitation Processes in the Eastern Ebro Subbasin (WISE-PreP) Project, Joan Bech et al.







CIFNCIA F INNOVACIÓN

